



STAR JPSS Annual Meeting
NOAA Center for Weather and Climate Prediction
College Park, MD



Naval Oceanographic Office (NAVOCEANO): Thermal uniformity fields and frontal regions

Jean-François Cayula

Vencore, Inc., Stennis Space Center, USA

Session 10: Sea Surface Temperature

August 11, 2016


Acknowledgments: This work was partially supported by the JPSS Program Office through a grant to the Northern Gulf Littoral Initiative and the University of South Mississippi. This study made extensive use of data from NOAA Comprehensive Large Array-data Stewardship System and GHRSSST Global Data Assembly Center at the NASA JPL PO.DAAC. Thanks to Doug May and Bruce Mckenzie of NAVOCEANO for the helpful discussions and encouragements. The views and findings contained in this presentation are those of the author and should not be construed as an official position, policy, or decision of NAVOCEANO or Vencore.

SST Uniformity field


- A Thermal Uniformity test is included in NAVOCEANO SST Processing.
- Uniformity field is defined as difference between min/max temperatures in a 3x3 pixels sliding window.
- A strict Thermal Uniformity test avoids contamination (fractional clouds) but also partially removes fronts.

SST Uniformity field

Example: SST Uniformity fields on January 16, 2016
for the Gulf of Mexico




NAVOCEANO SST Uniformity



ACSPO SST Uniformity. Note that in this example, ACSPO destriping step reduces the intrinsic noise.


SST Uniformity field

- Global SST uniformity fields show pattern corresponding to VIIRS aggregation scheme.



NAVOCEANO

July 14, 2014



ACSPO

Revised Uniformity Test

Recovery of Frontal Region

- Thermal Uniformity test on 3x3 pixel neighborhoods with 0.4°K threshold

•

When $T_{3\max} - T_{3\min} > 0.4^{\circ}\text{K}$

with $T_{3\max}/T_{3\min}$ being the maximum/minimum temperature
in the 3x3 pixel window

•


- ➔ Coherence of thermal gradient vector field
- ➔ Correlation between temperature and reflectance field

Revised Uniformity Test


Coherence of thermal gradient vector field

$$\text{coherence} = \frac{\| \text{mean}_w(\vec{\text{grad}}(x,y)) \|}{\| \text{mean}_w(| \vec{\text{grad}}(x,y) |) \|} < f \Rightarrow \text{not a front (cloud)}$$

- In noise free environment, f can be selected as a constant.
- Synthetic fields show the dependence of f on front strength. The
 - Step sizes vary from 0.3°K to 4°K
 - Standard deviation of noise is set 0.05°K
 - Probability of correct front detection is set at 95 percent



Example of synthetic field




f as function of front strength


Revised Uniformity Test

Coherence of thermal gradient vector field

- Result of using fixed and variable thresholds



Fixed threshold (0.5): Frontal regions that passed are in magenta.



Variable threshold: Frontal regions that passed are in color.

Revised Uniformity Test

Correlation of the thermal and reflective fields

- Performed when $T_{3\max} - T_{3\min} > 0.4^{\circ}\text{K}$
- Correlation between:
 - brightness temperature at $10.763\mu\text{m}$ (VIIRS M15) - bt
 - reflectance at $0.865\mu\text{m}$ (VIIRS M7) - r

$$\text{corr}(bt, r) = \frac{\text{cov}(bt, r)}{\text{stdev}(bt) * \text{stdev}(r)}$$


- Correlation is based on 7×7 pixel windows
- Negative correlation indicates cloud contamination leads to following decision rule:

$$\text{corr}(bt, r) < -0.6 \Rightarrow \text{not a front (cloud)}$$

Revised Uniformity Test


SST versus Brightness temperature

- Thermal uniformity and associated tests operate on Brightness Temperature. Using SST produces noisier results:



Correlation of the brightness temperature and reflectance fields.

Orbital overlap std=0.34°K



Correlation of the SST and reflectance fields.

Orbital overlap std=0.37°K


Revisited SST calculations

- The daytime NLSST or MCSST equations have the form:

$$\text{SST} = a * T_{11} + b * (T_{11} - T_{12}) \text{ where,}$$

$a \sim 1$ and $b \sim 2.5$ are semi constant variables

T_{11} and T_{12} are the brightness temperatures at $11\mu\text{m}$ and $12\mu\text{m}$




Daytime SST field on July 14, 2014

Revisited SST calculations


- SST uniformity field highlights noise in SST data.
- Noise which includes random noise and striping also corresponds to the VIIRS aggregation scheme.
- Much higher than noise of T_{11} brightness temperature data.

daytime Gulf Stream - composite sstminmax field




SST Uniformity field

daytime Gulf Stream - composite sstminmax field




T_{11} Uniformity field



Revisited SST calculations

- Revisited SST uses an $n \times n$ pixel average of the correction term on clean brightness temperature data.
- The resulting uniformity field is similar to that of the T_{11} field.
- Keeps SST front strength to at least T_{11} level
- Reduces effects of random noise and striping

Example
with $n=7$







Daytime revisited SST Uniformity field on July 14, 2014

Comparative results

Case 1: East coast of Japan

January 16, 2016, 140E to 143E and 35N to 37N

- Updated NAVOCEANO SST processing



SST field


SST uniformity field

Comparative results



Case 1: East coast of Japan

January 16, 2016, 140E to 143E and 35N to 37N

- Original NAVOCEANO SST processing



SST field



SST uniformity field




Comparative results



Case 1: East coast of Japan

January 16, 2016, 140E to 143E and 35N to 37N


- NOAA ACSPO SST processing (quality level 5 only)



SST field



SST uniformity field




Comparative results

Case 2: East coast of Argentina

January 16, 2016, 70W to 60W and 53S to 43S

- SST fields



Updated NAVOCEANO

NOAA ACSPO


Original NAVOCEANO

Comparative results

Case 2: East coast of Argentina

January 16, 2016, 70W to 60W and 53S to 43S

- SST Uniformity fields



Updated NAVOCEANO

NOAA ACSPO


Original NAVOCEANO

Comparative results

Case 2: East coast of Argentina

January 16, 2016, 70W to 60W and 53S to 43S

- Orbital overlap: SST differences between 2 consecutive orbits.
- Warm and cold spots often indicate contaminated data in one orbit



Updated NAVOCEANO
Std=0.24°K

NOAA ACSPO
Std=0.34°K

Original NAVOCEANO
Std=0.28°K

Global results

Orbital overlap evaluation

- January 16, 2016 from 120E to 30W* and 55S to 55N
- Standard deviation from orbital overlap evaluation

method	Number of retrievals	Standard deviation
Updated	61 millions	0.43°K
Original	68 millions	0.40°K
ACSPO	75 millions	0.50°K

Buoy match-ups

- Evaluation through buoy match-ups is preliminary. Early results indicate a standard deviation of 0.45°K for the updated and original NAVOCEANO SST (all categories), and 0.47°K for ACSPO SST (quality level 5 only).

*ACSPO SST missing data between 30W and 0W

Conclusion

- The updated NAVOCEANO SST processing successfully improves the coverage of frontal regions while maintaining strong cloud detection.
- The uniformity and associated tests perform better with brightness temperature than SST. This may have implications for SST edge detection.
- Replacing the standard correction term in the daytime SST equations by an $n \times n$ pixel average can drastically reduce the effect of random noise and striping while keeping the strength of the fronts in the resulting SST field to at least that of the level of the fronts in the brightness temperature field.




Comparative results

Case 3: Gulf of Mexico

January 16, 2016, 98W to 88W and 22N to 30N

- Updated NAVOCEANO SST processing



SST field


SST uniformity field

Comparative results

Case 3: Gulf of Mexico

January 16, 2016, 98W to 88W and 22N to 30N

- NOAA ACSPO SST processing



SST field


SST uniformity field

Comparative results

Case 3: Gulf of Mexico

January 16, 2016, 98W to 88W and 22N to 30N

- Original NAVOCEANO SST processing



SST field

SST uniformity field